arXiv:1408.4400v1 [math.FA] 19 Aug 2014

Maximally Modulated Singular Integral Operators and their Applications to Pseudodifferential Operators on Banach Function Spaces

Alexei Yu. Karlovich

Abstract. We prove that if the Hardy-Littlewood maximal operator is bounded on a separable Banach function space $X(\mathbb{R}^n)$ and on its associate space $X'(\mathbb{R}^n)$ and a maximally modulated Calderón-Zygmund singular integral operator T^Φ is of weak type (r,r) for all $r\in(1,\infty)$, then T^Φ extends to a bounded operator on $X(\mathbb{R}^n)$. This theorem implies the boundedness of the maximally modulated Hilbert transform on variable Lebesgue spaces $L^{p(\cdot)}(\mathbb{R})$ under natural assumptions on the variable exponent $p:\mathbb{R}\to(1,\infty)$. Applications of the above result to the boundedness and compactness of pseudodifferential operators with $L^\infty(\mathbb{R},V(\mathbb{R}))$-symbols on variable Lebesgue spaces $L^{p(\cdot)}(\mathbb{R})$ are considered. Here the Banach algebra $L^\infty(\mathbb{R},V(\mathbb{R}))$ consists of all bounded measurable $V(\mathbb{R})$-valued functions on $\mathbb{R}$ where $V(\mathbb{R})$ is the Banach algebra of all functions of bounded total variation.

1. Introduction

In this paper we will be concerned with the boundedness of maximally modulated Calderón-Zygmund singular integral operators and its applications to the boundedness of pseudodifferential operators with non-regular symbols on separable Banach function spaces.

Let us define the main operators we are dealing with. Let $L_0^\infty(\mathbb{R}^n)$ and $C_0^\infty(\mathbb{R}^n)$ denote the sets of all bounded functions with compact support and all infinitely differentiable functions with compact support, respectively. A Calderón-Zygmund operator is a linear operator T which is bounded on $L^2(\mathbb{R}^n)$ such that for every $f\in L_0^\infty(\mathbb{R}^n)$,

$$(Tf)(x):=\int_{\mathbb{R}^n}K(x,y)f(y)\,dy\quad\text{for a.e.}\quad x\in\mathbb{R}^n\setminus\operatorname{supp}f,$$

2010 *Mathematics Subject Classification.* Primary 42B20, 47G30; Secondary 46E30, 42B25.

Key words and phrases. Maximally modulated singular integral operator, Calderón-Zygmund operator, Hilbert transform, pseudodifferential operator with non-regular symbol, Banach function space, variable Lebesgue space.

This work was partially supported by the Fundação para a Ciência e a Tecnologia (Portuguese Foundation for Science and Technology) through the projects PEst-OE/MAT/UI0297/2014 (Centro de Matemática e Aplicações) and EXPL/MAT-CAL/0840/2013 (Problemas Variacionais em Espaços de Sobolev de Expoente Variável).

where $\operatorname{supp} f$ denotes the support of f. The kernel

$$K : \mathbb{R}^n \times \mathbb{R}^n \setminus \{(x,x) : x \in \mathbb{R}^n\} \to \mathbb{C}$$

is assumed to satisfy the following standard conditions:

$$|K(x,y)| \le \frac{c_0}{|x-y|^n} \quad \text{for} \quad x \ne y$$

and

$$|K(x,y) - K(x,y')| + |K(y,x) - K(y',x)| \le \frac{c_0 |y-y'|^\tau}{|x-y|^{n+\tau}}$$

for $|x-y| > 2|y-y'|$, where c_0 and τ are some positive constants independent of $x, y, y' \in \mathbb{R}^n$ (see, e.g., [**G09**, Section 8.1.1]). The most prominent example of Calderón-Zygmund operators is the Hilbert transform defined for $f \in L_0^\infty(\mathbb{R})$ by

$$(Hf)(x) := \frac{1}{\pi} \text{v.p.} \int_{\mathbb{R}} \frac{f(y)}{x-y}\,dy = \lim_{\varepsilon \to 0} \frac{1}{\pi} \int_{\mathbb{R} \setminus I(x,\varepsilon)} \frac{f(y)}{x-y}\,dy, \quad x \in \mathbb{R},$$

where $I(x,\varepsilon) := (x-\varepsilon, x+\varepsilon)$.

Suppose $\Phi = \{\phi_\alpha\}_{\alpha \in \mathcal{A}}$ is a family of measurable real-valued functions indexed by an arbitrary set $\mathcal{A}$. Then for every $\phi_\alpha \in \mathcal{A}$, the modulation operator is defined by

$$(\mathcal{M}^{\phi_\alpha} f)(x) := e^{-i\phi_\alpha(x)} f(x), \quad x \in \mathbb{R}^n.$$

Following [**GMS05**] (see also [**DPL13**]), the maximally modulated singular integral operator T^Φ of the Calderón-Zygmund operator T with respect to the family Φ is defined for $f \in L_0^\infty(\mathbb{R}^n)$ by

$$(T^\Phi f)(x) := \sup_{\alpha \in \mathcal{A}} |T(\mathcal{M}^{\phi_\alpha} f)(x)|, \quad x \in \mathbb{R}^n.$$

This definition is motivated by the fact that the maximally modulated Hilbert transform

$$(\mathcal{C}f)(x) := (H^\Psi f)(x) \quad \text{with} \quad \Psi := \{\psi_\alpha(x) = \alpha x : \alpha, x \in \mathbb{R}\}$$

is closely related to the continuous version of the celebrated Carleson-Hunt theorem on the a.e. convergence of Fourier series (see, e.g., [**D91**, Chap. 2, Section 2.2], [**G09**, Chap. 11], and [**MS13**, Chap. 7]). In [**GMS05, DPL13**] the operator $\mathcal{C}$ is called the Carleson operator, however in [**G09**, Section 11.1] and in [**MS13**, Section 7.1] this term is used for two different from $\mathcal{C}$ and each other operators.

For $f \in C_0^\infty(\mathbb{R})$, consider the maximal singular integral operator given by

$$(S_* f)(x) := \sup_{-\infty < a < b < +\infty} \left| (S_{(a,b)} f)(x) \right|, \quad x \in \mathbb{R}, \tag{1.1}$$

where $S_{(a,b)} f$ is the integral analogue of the partial sum of the Fourier series given by

$$(S_{(a,b)} f)(x) := \frac{1}{2\pi} \int_a^b \widehat{f}(\lambda) e^{ix\lambda}\,d\lambda, \quad x \in \mathbb{R},$$

and

$$\widehat{f}(\lambda) := (\mathcal{F}f)(\lambda) := \int_{\mathbb{R}} f(x) e^{-ix\lambda} dx, \quad \lambda \in \mathbb{R},$$

is the Fourier transform of f. It is not difficult to see that if $f \in C_0^\infty(\mathbb{R})$, then

$$(S_* f)(x) \le (\mathcal{C}f)(x) \quad \text{for a.e.} \quad x \in \mathbb{R}. \tag{1.2}$$

For a suitable function a on $\mathbb{R}\times\mathbb{R}$, a pseudodiffferential operator $a(x,D)$ is defined for a function $f\in C_0^\infty(\mathbb{R})$ by the iterated integral

$$(a(x,D)f)(x) := \frac{1}{2\pi}\int_{\mathbb{R}} d\lambda \int_{\mathbb{R}} a(x,\lambda)e^{i(x-y)\lambda}f(y)\,dy, \quad x\in\mathbb{R}. \tag{1.3}$$

The function a is called the symbol of the pseudodifferential operator $a(x,D)$.

Our results on the above mentioned operators will be formulated in terms of the Hardy-Littlewood maximal function, which we define next. Let $1\le r<\infty$. Given $f\in L^r_{\mathrm{loc}}(\mathbb{R}^n)$, the r-th maximal operator is defined by

$$(M_r f)(x) := \sup_{Q\ni x}\left(\frac{1}{|Q|}\int_Q |f(y)|^r dy\right)^{1/r}, \quad x\in\mathbb{R}^n,$$

where the supremum is taken over all cubes Q containing x. Here, and throughout, all cubes will be assumed to have their sides parallel to the coordinate axes and $|Q|$ will denote the volume of Q. For $r=1$ this is the usual Hardy-Littlewood maximal operator, which will be denoted by M.

Let $f\in L^1_{\mathrm{loc}}(\mathbb{R}^n)$. For a cube $Q\subset\mathbb{R}^n$, put

$$f_Q := \frac{1}{|Q|}\int_Q f(x)dx.$$

The Fefferman-Stein sharp maximal operator $f\mapsto M^\# f$ is defined by

$$(M^\# f)(x) := \sup_{Q\ni x}\frac{1}{|Q|}\int_Q |f(x)-f_Q|dx, \quad x\in\mathbb{R}^n,$$

where the supremum is taken over all cubes Q containing x.

Banach function spaces $X(\mathbb{R}^n)$ will be defined in Section 2. This is a wide class of spaces including rearrangement-invariant (r.i.) Lebesgue, Orlicz, and Lorentz spaces, as well as non-r.i. variable Lebesgue spaces $L^{p(\cdot)}(\mathbb{R}^n)$. The main feature of these spaces is the so-called lattice property: if $|f(x)|\le|g(x)|$ for a.e. $x\in\mathbb{R}^n$, then $\|f\|_{X(\mathbb{R}^n)}\le\|g\|_{X(\mathbb{R}^n)}$. In Section 2 we collect preliminaries prepare the proof of main results given in Section 3. Let us briefly describe them.

The boundedness of maximally modulated Calderón-Zygmund operators T^Φ on weighted Lebesgue spaces was studied by Grafakos, Martell, and Soria [**GMS05**]. A quantitative version of their results was obtained recently by Di Plinio and Lerner [**DPL13**]. We show that a pointwise inequality for the sharp maximal function of T^Φ obtained in [**GMS05**, Proposition 4.1], combined with the Fefferman-Stein inequality for Banach function spaces due to Lerner [**L10**, Corollary 4.2], and with the self-improving property of the Hardy-Littlewood maximal function on Banach function spaces obtained by Lerner and Pérez [**LP07**, Corolary 1.3], imply the boundedness of T^Φ on a separable Banach function space $X(\mathbb{R}^n)$ under the natural assumptions that M is bounded on $X(\mathbb{R}^n)$, M is bounded on its associate space $X'(\mathbb{R}^n)$, and T^Φ is of weak type (r,r) for all $r\in(1,\infty)$ (see Theorem 3.1). Notice that the latter hypothesis is satisfied for the maximally modulated Hilbert transform $\mathcal{C}$. This gives the boundedness of $\mathcal{C}$ on $X(\mathbb{R})$ (see Corollary 3.3). From here and the pointwise estimate (1.2) we also get the boundedness of the operator S_* on separable Banach functions spaces such that M is bounded on $X(\mathbb{R})$ and on $X'(\mathbb{R})$ (see Lemma 4.2).

Section 4 is devoted to applications of the above results to the boundedness of pseudodifferential operators with non-regular symbols on Banach function

spaces. Note that the boundedness of $a(x,D)$ with smooth (regular) symbols in Hörmander's and Miyachi's classes on separable Banach function spaces was studied in [**K-A14**] (see also [**KS13**]). On the other hand, Yu. Karlovich [**K-Yu07**] introduced the class $L^\infty(\mathbb{R},V(\mathbb{R}))$ of bounded measurable $V(\mathbb{R})$-valued functions on $\mathbb{R}$ where $V(\mathbb{R})$ is the Banach algebra of all functions of bounded total variation. Symbols in $L^\infty(\mathbb{R},V(\mathbb{R}))$ may have jump discontinuities in both variables. By using the boundedness of the operator S_* on $L^p(\mathbb{R})$ for $1<p<\infty$, Yu. Karlovich [**K-Yu07**, Theorem 3.1] proved the boundedness of $a(x,D)$ with $a\in L^\infty(\mathbb{R},V(\mathbb{R}))$ on $L^p(\mathbb{R})$ for $1<p<\infty$. Later on he extended this result to weighted Lebesgue spaces $L^p(\mathbb{R},w)$ with Muckenhoupt weights w (see [**K-Yu12**, Theorem 4.1]). One of the important ingredients of those proofs is the pointwise inequality

$$|(a(x,D)f)(x)|\le 2(S_*f)(x)\|a(x,\cdot)\|_V,\quad x\in\mathbb{R},$$

for $f\in C_0^\infty(\mathbb{R})$. From this inequality and the boundedness of S_* we obtain the boundedness of $a(x,D)$ on separable Banach function spaces $X(\mathbb{R})$ under the assumption that M is bounded on $X(\mathbb{R})$ and $X'(\mathbb{R})$ (see Theorem 4.3).

In Section 5 we specify the above results to the case of variable Lebesgue spaces $L^{p(\cdot)}(\mathbb{R}^n)$. If the variable exponent $p:\mathbb{R}^n\to[1,\infty]$ is bounded away from one and infinity, then in view of Diening's theorem [**D05**, Theorem 8.1], the boundedness of M is equivalent to the boundedness of M on its associate space. Hence all above results have simpler formulations in the case of variable Lebesgue spaces (see Section 5).

We conclude the paper with a sufficient condition for the compactness of pseudodifferential operators $a(x,D)$ with $L^\infty(\mathbb{R},V(\mathbb{R}))$-symbols on variable Lebesgue spaces $L^{p(\cdot)}(\mathbb{R})$ (see Corollary 6.5). This result is obtained from Yu. Karlovich's result [**K-Yu07**, Theorem 4.1] for standard Lebesgue spaces by transferring the compactness property from standard to variable Lebesgue spaces with the aid of the Krasnosel'skii-type interpolation theorem (see Section 6.2).

2. Preliminaries

2.1. Banach function spaces. The set of all Lebesgue measurable complex-valued functions on $\mathbb{R}^n$ is denoted by $\mathcal{M}$. Let $\mathcal{M}^+$ be the subset of functions in $\mathcal{M}$ whose values lie in $[0,\infty]$. The characteristic function of a measurable set $E\subset\mathbb{R}^n$ is denoted by χ_E and the Lebesgue measure of E is denoted by $|E|$.

Definition 2.1 ([**BS88**, Chap. 1, Definition 1.1]). A mapping $\rho:\mathcal{M}^+\to[0,\infty]$ is called a *Banach function norm* if, for all functions f,g,f_n $(n\in\mathbb{N})$ in $\mathcal{M}^+$, for all constants $a\ge 0$, and for all measurable subsets E of $\mathbb{R}^n$, the following properties hold:

(A1) $\rho(f)=0\Leftrightarrow f=0$ a.e., $\quad\rho(af)=a\rho(f)$, $\quad\rho(f+g)\le\rho(f)+\rho(g)$,

(A2) $0\le g\le f$ a.e. $\Rightarrow\ \rho(g)\le\rho(f)$ (the lattice property),

(A3) $0\le f_n\uparrow f$ a.e. $\Rightarrow\ \rho(f_n)\uparrow\rho(f)$ (the Fatou property),

(A4) $|E|<\infty\Rightarrow\rho(\chi_E)<\infty$,

(A5) $|E|<\infty\Rightarrow\displaystyle\int_E f(x)\,dx\le C_E\rho(f)$

with $C_E\in(0,\infty)$ which may depend on E and ρ but is independent of f.

When functions differing only on a set of measure zero are identified, the set $X(\mathbb{R}^n)$ of all functions $f \in \mathcal{M}$ for which $\rho(|f|) < \infty$ is called a *Banach function space*. For each $f \in X(\mathbb{R}^n)$, the norm of f is defined by

$$\|f\|_{X(\mathbb{R}^n)} := \rho(|f|).$$

The set $X(\mathbb{R}^n)$ under the natural linear space operations and under this norm becomes a Banach space (see [**BS88**, Chap. 1, Theorems 1.4 and 1.6]).

The norm of a bounded sublinear operator A on a Banach function space $X(\mathbb{R}^n)$ will be denoted by $\|A\|_{\mathcal{B}(X(\mathbb{R}^n))}$.

If ρ is a Banach function norm, its associate norm ρ' is defined on $\mathcal{M}^+$ by

$$\rho'(g) := \sup\left\{\int_{\mathbb{R}^n} f(x)g(x)\,dx \ : \ f \in \mathcal{M}^+,\ \rho(f) \le 1\right\}, \quad g \in \mathcal{M}^+.$$

It is a Banach function norm itself [**BS88**, Chap. 1, Theorem 2.2]. The Banach function space $X'(\mathbb{R}^n)$ determined by the Banach function norm ρ' is called the *associate space* (*Köthe dual*) of $X(\mathbb{R}^n)$. The Lebesgue space $L^p(\mathbb{R}^n)$, $1 \le p \le \infty$, are the archetypical example of Banach function spaces. Other classical examples of Banach function spaces are Orlicz spaces, rearrangement-invariant spaces, and variable Lebesgue spaces $L^{p(\cdot)}(\mathbb{R}^n)$.

2.2. Density of bounded and smooth compactly supported functions in separable Banach function spaces. The proof of the following fact is standard. For details, see [**KS14**, Lemma 2.10(b)], where it was proved for $n = 1$. The proof for arbitrary n is a minor modification of that one.

LEMMA 2.2. *The sets $L_0^\infty(\mathbb{R}^n)$ and $C_0^\infty(\mathbb{R}^n)$ are dense in a separable Banach function space $X(\mathbb{R}^n)$.*

2.3. Nonnegative sublinear operators on Banach function spaces. The operators T^Φ, $\mathcal{C}$, and $S_{(a,b)}$ although nonlinear, are examples of sublinear operators that assume only nonnegative values. Let us give a precise definition of this class of operators. Let D be a linear subspace of $\mathcal{M}$. An operator $T : D \to \mathcal{M}$ is said to be nonnegative sublinear (cf. [**BS88**, p. 230]) if

$$0 \le T(f+g) \le Tf + Tg, \quad T(\lambda f) = |\lambda| Tf \quad \text{a.e. on } \mathbb{R}^n \tag{2.1}$$

for all $f, g \in D$ and all constants $\lambda \in \mathbb{C}$. The following result is well known for linear operators. With the property

$$|Tf - Tg| \le |T(f-g)| = T(f-g), \quad f, g \in D,$$

which is an immediate consequence of (2.1), essentially the same proof establishes the result also for nonnegative sublinear operators.

LEMMA 2.3. *Let D be a dense linear subspace of a Banach function space $X(\mathbb{R}^n)$ and $T : D \to \mathcal{M}$ be a nonnegative sublinear operator. If there exists a positive constant C such that*

$$\|Tf\|_{X(\mathbb{R}^n)} \le C\|f\|_{X(\mathbb{R}^n)} \quad \textit{for all} \quad f \in D,$$

then T has a unique extension to a nonnegative sublinear operator $\widetilde{T} : X(\mathbb{R}^n) \to \mathcal{M}$ such that

$$\|\widetilde{T}f\|_{X(\mathbb{R}^n)} \le C\|f\|_{X(\mathbb{R}^n)} \quad \textit{for all} \quad f \in X(\mathbb{R}^n).$$

In what follows we will use the same notation for an operator defined on a dense subspace and for its bounded extension to the whole space.

2.4. Self-improving property of maximal operators on Banach function spaces. If $1 < q < \infty$, then from the Hölder inequality one can immediately get that

$$(Mf)(x) \le (M_r f)(x) \qquad \text{for a.e.} \quad x \in \mathbb{R}^n.$$

Thus, the boundedness of any M_r, $1 < r < \infty$, on a Banach function space $X(\mathbb{R}^n)$ immediately implies the boundedness of M. A partial converse of this fact, called a *self-improving property* of the Hardy-Littlewood maximal operator, is also true. It was proved by Lerner and Pérez [**LP07**] (see also [**LO10**] for another proof) in a more general setting of quasi-Banach function spaces.

THEOREM 2.4 ([**LP07**, Corollary 1.3]). *Let $X(\mathbb{R}^n)$ be a Banach function space. Then M is bounded on $X(\mathbb{R}^n)$ if and only if M_r is bounded on $X(\mathbb{R}^n)$ for some $r \in (1, \infty)$.*

2.5. The Fefferman-Stein inequality for Banach function spaces. It is obvious that $M^\# f$ is pointwise dominated by Mf. Hence, by Axiom (A2),

$$\|M^\# f\|_{X(\mathbb{R}^n)} \le \text{const} \|f\|_{X(\mathbb{R}^n)} \quad \text{for} \quad f \in X(\mathbb{R}^n)$$

whenever M is bounded on $X(\mathbb{R}^n)$. The converse inequality for Lebesgue spaces $L^p(\mathbb{R}^n)$, $1 < p < \infty$, was proved by Fefferman and Stein (see [**FS72**, Theorem 5] and also [**S93**, Chap. IV, Section 2.2]). The following extension of the Fefferman-Stein inequality to Banach function spaces was proved by Lerner [**L10**].

Let $S_0(\mathbb{R}^n)$ be the space of all measurable functions f on $\mathbb{R}^n$ such that

$$|\{x \in \mathbb{R}^n : |f(x)| > \lambda\}| < \infty \quad \text{for all} \quad \lambda > 0.$$

THEOREM 2.5 ([**L10**, Corollary 4.2]). *Let M be bounded on a Banach function space $X(\mathbb{R}^n)$. Then M is bounded on its associate space $X'(\mathbb{R}^n)$ if and only if there exists a constant $C_\# > 0$ such that, for all $f \in S_0(\mathbb{R}^n)$,*

$$\|f\|_{X(\mathbb{R}^n)} \le C_\# \|M^\# f\|_{X(\mathbb{R}^n)}.$$

2.6. Pointwise inequality for the sharp maximal function of T^Φ. Let $1 \le r < \infty$. Recall that a sublinear operator $A : L^r(\mathbb{R}^n) \to \mathcal{M}$ is said to be of weak type (r, r) if

$$|\{x \in \mathbb{R}^n : |(Af)(x)| > \lambda\}| \le \frac{C^r}{\lambda^r} \int_{\mathbb{R}^n} |f(y)|^r \, dy$$

for all $f \in L^r(\mathbb{R}^n)$ and $\lambda > 0$, where C is a positive constant independent of f and λ. It is well known that if A is bounded on the standard Lebesgue space $L^r(\mathbb{R}^n)$, then it is of weak type (r, r).

Grafakos, Martell, and Soria [**GMS05**] developed two alternative approaches to weighted L^r estimates for maximally modulated Calderón-Zygmund singular integral operators T^Φ. One is based on good-λ inequalities, another rests on the following pointwise estimate for the sharp maximal function of T^Φ.

LEMMA 2.6 ([**GMS05**, Proposition 4.1]). *Suppose T is a Calderón-Zygmund operator and $\Phi = \{\phi_\alpha\}_{\alpha \in \mathcal{A}}$ is a family of measurable real-valued functions indexed by an arbitrary set $\mathcal{A}$. If T^Φ is of weak type (r, r) for some $r \in (1, \infty)$, then there is a positive constant C_r such that for every $f \in L_0^\infty(\mathbb{R}^n)$,*

$$M^\#(T^\Phi f)(x) \le C_r M_r f(x) \quad \text{for a.e.} \quad x \in \mathbb{R}^n.$$

3. Maximally modulated singular integrals on Banach function spaces

3.1. Boundedness of maximally modulated Calderón-Zygmund singular integral operators on Banach function spaces. We are in a position to prove the main result of the paper.

THEOREM 3.1. *Let $X(\mathbb{R}^n)$ be a separable Banach function space. Suppose the Hardy-Littlewood maximal operator M is bounded on $X(\mathbb{R}^n)$ and on its associate space $X'(\mathbb{R}^n)$. Suppose T is a Calderón-Zygmund operator and $\Phi = \{\phi_\alpha\}_{\alpha\in\mathcal{A}}$ is a family of measurable real-valued functions indexed by an arbitrary set $\mathcal{A}$. If T^Φ is of weak type (r,r) for all $r \in (1,\infty)$, then T^Φ extends to a bounded operator on the space $X(\mathbb{R}^n)$.*

PROOF. We argue as in the proof of [**K-A14**, Theorem 1.2]. Since M is bounded on $X(\mathbb{R}^n)$, by Theorem 2.4, there is an $r \in (1,\infty)$ such that the maximal function M_r is bounded on $X(\mathbb{R}^n)$, that is, there is a positive constant C such that

$$\|(M_r\varphi)\|_{X(\mathbb{R}^n)} \le C\|\varphi\|_{X(\mathbb{R}^n)} \quad \text{for all} \quad \varphi \in X(\mathbb{R}^n). \tag{3.1}$$

Assume that $f \in C_0^\infty(\mathbb{R}^n)$. By the hypothesis, T^Φ is of weak type (r,r) and M is bounded on $X'(\mathbb{R}^n)$. Therefore, $T^\Phi f \in S_0(\mathbb{R}^n)$. Moreover, by Theorem 2.5, there exists a positive constant $C_\#$ such that

$$\|T^\Phi f\|_{X(\mathbb{R}^n)} \le C_\#\|M^\#(T^\Phi f)\|_{X(\mathbb{R}^n)} \quad \text{for all} \quad f \in C_0^\infty(\mathbb{R}^n). \tag{3.2}$$

From Lemma 2.6 and Axioms (A1)–(A2) we conclude that there exists a positive constant C_r such that

$$\|M^\#(T^\Phi f)\|_{X(\mathbb{R}^n)} \le C_r\|M_r f\|_{X(\mathbb{R}^n)} \quad \text{for all} \quad f \in C_0^\infty(\mathbb{R}^n). \tag{3.3}$$

Combining inequalities (3.1)–(3.3), we arrive at

$$\|T^\Phi f\|_{X(\mathbb{R}^n)} \le CC_\#C_r\|f\|_{X(\mathbb{R}^n)} \quad \text{for all} \quad f \in C_0^\infty(\mathbb{R}^n).$$

To conclude the proof, it remains to recall that $C_0^\infty(\mathbb{R}^n)$ is dense in the separable Banach function space $X(\mathbb{R}^n)$ in view of Lemma 2.2 and apply Lemma 2.3. □

3.2. Boundedness of the maximally modulated Hilbert transform on standard Lebesgue spaces. Fix $f \in L^1_{loc}(\mathbb{R})$. Let H_* be the maximal Hilbert transform given by

$$(H_*f)(x) := \sup_{\varepsilon>0}\left|\frac{1}{\pi}\int_{\mathbb{R}\setminus I(x,\varepsilon)}\frac{f(y)}{x-y}\,dy\right|,$$

where $I(x,\varepsilon) = (x-\varepsilon, x+\varepsilon)$. Further, let $\mathcal{C}_*$ be the maximally modulated maximal Hilbert transform (called also the maximal Carleson operator) defined by

$$(\mathcal{C}_*f)(x) := \sup_{a\in\mathbb{R}}(H_*(\mathcal{M}^{\psi_a}f))(x) \quad \text{with} \quad \psi_a(x) = ax, \quad a,x \in \mathbb{R}.$$

It is easy to see that

$$(\mathcal{C}f)(x) \le (\mathcal{C}_*f)(x), \quad x \in \mathbb{R}. \tag{3.4}$$

The boundedness of the operator $\mathcal{C}_*$ on the standard Lebesgue spaces $L^r(\mathbb{R})$ is proved, e.g., in [**G09**, Theorem 11.3.3] (see also [**K-Yu12**, Theorem 2.7]). From this observation and (3.4) we get the following result (see also [**D91**, Theorem 2.1] and [**K-Yu12**, Theorem 2.8]).

LEMMA 3.2. *The maximally modulated Hilbert transform $\mathcal{C}$ is bounded on every standard Lebesgue space $L^r(\mathbb{R})$ for $1 < r < \infty$.*

3.3. Boundedness of the maximally modulated Hilbert transform on separable Banach function spaces. From Theorem 3.1 and Lemma 3.2 we immediately get the following.

COROLLARY 3.3. *Let $X(\mathbb{R})$ be a separable Banach function space. Suppose the Hardy-Littlewood maximal operator M is bounded on $X(\mathbb{R})$ and on its associate space $X'(\mathbb{R})$. Then the maximally modulated Hilbert transform $\mathcal{C}$ extends to a bounded operator on $X(\mathbb{R})$.*

4. Boundedness of pseudodifferential operators with non-regular symbols on Banach function spaces

4.1. Functions of bounded total variation. Let a be a complex-valued function of bounded total variation $V(a)$ on $\mathbb{R}$ where

$$V(a) := \sup\left\{\sum_{k=1}^{n} |a(x_k) - a(x_{k-1})| : -\infty < x_0 < x_1 < \cdots < x_n < +\infty, n \in \mathbb{N}\right\}$$

Hence at every point $x \in \dot{\mathbb{R}} := \mathbb{R} \cup \{\infty\}$ the one-sided limits

$$a(x \pm 0) = \lim_{t \to x^{\pm}} a(t)$$

exist, where $a(\pm\infty) = a(\infty \mp 0)$, and the set of discontinuities of a is at most countable (see, e.g., [**N55**, Chap. VIII, Sections 3 and 9]). Without loss of generality we will assume that functions of bounded total variation are continuous from the left at every discontinuity point $x \in \dot{R}$. The set $V(\mathbb{R})$ of all continuous from the left functions of bounded total variation on $\mathbb{R}$ is a unital non-separable Banach algebra with the norm

$$\|a\|_V := \|a\|_{L^\infty(\mathbb{R})} + V(a).$$

By analogy with $V(a) = V_{-\infty}^{+\infty}(a)$, one can define the total variations $V_c^d(a)$, $V_{-\infty}^c(a)$, and $V_d^{+\infty}(a)$ of a function $a : \mathbb{R} \to \mathbb{C}$ on $[c, d]$, $(-\infty, c]$, and $[d, +\infty)$, taking, respectively, the partitions

$$c = x_0 < x_1 < \cdots < x_n = d, \quad -\infty < x_0 < x_1 < \cdots < x_n = c,$$

and $d = x_0 < x_1 < \cdots < x_n < +\infty$.

4.2. Non-regular symbols of pseudodifferential operators. Following [**K-Yu07, K-Yu12**], we denote by $L^\infty(\mathbb{R}, V(\mathbb{R}))$ the set of functions $a : \mathbb{R} \times \mathbb{R} \to \mathbb{C}$ such that $\widehat{a} : x \mapsto a(x, \cdot)$ is a bounded measurable $V(\mathbb{R})$-valued function on $\mathbb{R}$. Note that in view of non-separability of the Banach space $V(\mathbb{R})$, the measurability of $\widehat{a}$ means that the map $\widehat{a} : \mathbb{R} \to V(\mathbb{R})$ possesses the Luzin property: for any compact set $K \subset \mathbb{R}$ and any δ there is a compact set $K_\delta \subset K$ such that $|K \setminus K_\delta| < \delta$ and $\widehat{a}$ is continuous on K_δ (see, e.g., [**S67**, Chap. IV, Section 4, p. 487]). This implies that the function $x \mapsto a(x, \lambda \pm 0)$ for all $\lambda \in \dot{\mathbb{R}}$ and the function $x \mapsto \|a(x, \cdot)\|_V$ are measurable on $\mathbb{R}$ as well. Note that for almost all $x \in \mathbb{R}$ the limits $a(x, \lambda \pm 0)$ the limits exist for all $\lambda \in \dot{\mathbb{R}}$, $a(x, \lambda) = a(x, \lambda - 0)$ for all $\lambda \in \mathbb{R}$ and we put

$$a(x, \pm\infty) := \lim_{\lambda \to \pm\infty} a(x, \lambda).$$

Therefore, the functions $a(\cdot,\lambda\pm 0)$ for every $\lambda\in\dot{\mathbb{R}}$ and the function $x\mapsto\|a(x,\cdot)\|_V$, where

$$\|a(x,\cdot)\|_V := \|a(x,\cdot)\|_{L^\infty(\mathbb{R})} + V(a(x,\cdot)),$$

belong to $L^\infty(\mathbb{R})$. Clearly, $L^\infty(\mathbb{R},V(\mathbb{R}))$ is a unital Banach algebra with the norm

$$\|a\|_{L^\infty(\mathbb{R},V(\mathbb{R}))} = \operatorname*{ess\,sup}_{x\in\mathbb{R}} \|a(x,\cdot)\|_V.$$

4.3. Pointwise inequality for pseudodifferential operators. The following pointwise estimate was obtained by Yuri Karlovich in the proof of [**K-Yu07**, Theorem 3.1] and [**K-Yu12**, Theorem 4.1].

LEMMA 4.1. *If $a\in L^\infty(\mathbb{R},V(\mathbb{R}))$ and $f\in C_0^\infty(\mathbb{R})$, then*

$$\big|(a(x,D)f)(x)\big| \le 2(S_*f)(x)\|a(x,\cdot)\|_V \quad \text{for a.e.} \quad x\in\mathbb{R}.$$

4.4. Boundedness of the maximal singular integral operator S_* on separable Banach function spaces. We continue with the following result on the boundedness of the maximal singular integral operator S_* initially defined for $f\in C_0^\infty(\mathbb{R})$ by (1.1).

LEMMA 4.2. *Let $X(\mathbb{R})$ be a separable Banach function space. Suppose the Hardy-Littlewood maximal operator M is bounded on $X(\mathbb{R})$ and on its associate space $X'(\mathbb{R})$. Then the operator S_*, defined for the functions $f\in C_0^\infty(\mathbb{R})$ by* (1.1)*, extends to a bounded operator on the space $X(\mathbb{R})$.*

PROOF. Fix $f\in C_0^\infty(\mathbb{R})$. It is not difficult to check (see, e.g., [**D91**, Chap. 2, Section 2.2] and also [**G09**, p. 475]) that

$$(S_{(a,b)}f)(x) = \frac{i}{2}\left\{\mathcal{M}^{-\psi_a}\big(H(\mathcal{M}^{\psi_a}f)\big)(x) - \mathcal{M}^{-\psi_b}\big(H(\mathcal{M}^{\psi_b}f)\big)(x)\right\}, \quad x\in\mathbb{R},$$

where $\psi_a(x)=ax$, $\psi_b(x)=bx$ and $-\infty<a<b<+\infty$. Therefore,

$$\begin{aligned}(S_*f)(x) &= \sup_{-\infty<a<b<\infty} |(S_{(a,b)}f)(x)| \\ &\le \frac{1}{2}\sup_{a\in\mathbb{R}}\big|\big(H(\mathcal{M}^{\psi_a}f)\big)(x)\big| + \frac{1}{2}\sup_{b\in\mathbb{R}}\big|\big(H(\mathcal{M}^{\psi_b}f)\big)(x)\big| \\ &= (\mathcal{C}f)(x), \quad x\in\mathbb{R}.\end{aligned}$$

From this inequality, Axioms (A1)-(A2), and Corollary 3.3 we get

$$\|S_*f\|_{X(\mathbb{R})} \le \|\mathcal{C}f\|_{X(\mathbb{R})} \le \|\mathcal{C}\|_{\mathcal{B}(X(\mathbb{R}))}\|f\|_{X(\mathbb{R})} \quad \text{for} \quad f\in C_0^\infty(\mathbb{R}).$$

It remains to apply Lemma 2.2. □

4.5. Boundedness of pseudodifferential operators with $L^\infty(\mathbb{R},V(\mathbb{R}))$ symbols on Banach function spaces. We are ready to prove the boundedness result for pseudodifferential operators with non-regular symbols on separable Banach function spaces.

THEOREM 4.3. *Let $X(\mathbb{R})$ be a separable Banach function space. Suppose the Hardy-Littlewood maximal operator M is bounded on $X(\mathbb{R})$ and on its associate space $X'(\mathbb{R})$. If $a\in L^\infty(\mathbb{R},V(\mathbb{R}))$, then the pseudodifferential operator $a(x,D)$, defined for the functions $f\in C_0^\infty(\mathbb{R})$ by the iterated integral* (1.3)*, extends to a bounded linear operator on the space $X(\mathbb{R})$ and*

$$\|a(x,D)\|_{\mathcal{B}(X(\mathbb{R}))} \le 2\|S_*\|_{\mathcal{B}(X(\mathbb{R}))}\|a\|_{L^\infty(\mathbb{R},V(\mathbb{R}))}.$$

PROOF. From Lemma 4.1, axioms (A1)-(A2), and Lemma 4.2 we obtain for $f \in C_0^\infty(\mathbb{R})$,

$$\begin{aligned}\|a(x,D)f\|_{X(\mathbb{R})} &\le 2\|S_*f\|_{X(\mathbb{R})} \operatorname*{ess\,sup}_{x\in\mathbb{R}} \|a(x,\cdot)\|_V \\ &\le 2\|S_*\|_{\mathcal{B}(X(\mathbb{R}))}\|a\|_{L^\infty(\mathbb{R},V(\mathbb{R}))}\|f\|_{X(\mathbb{R})}.\end{aligned}$$

Since $C_0^\infty(\mathbb{R})$ is dense in the space $X(\mathbb{R})$ in view of Lemma 2.2, from the above estimate we arrive immediately at the desired conclusion. □

5. Boundedness of maximally modulated Calderón-Zygmund operators and pseudodifferential operators with non-regular symbols on variable Lebesgue spaces

5.1. Variable Lebesgue spaces. Let $p\colon \mathbb{R}^n \to [1,\infty]$ be a measurable a.e. finite function. By $L^{p(\cdot)}(\mathbb{R}^n)$ we denote the set of all complex-valued functions f on $\mathbb{R}$ such that

$$I_{p(\cdot)}(f/\lambda) := \int_{\mathbb{R}^n} |f(x)/\lambda|^{p(x)}dx < \infty$$

for some $\lambda > 0$. This set becomes a Banach function space when equipped with the norm

$$\|f\|_{p(\cdot)} := \inf\big\{\lambda > 0 : I_{p(\cdot)}(f/\lambda) \le 1\big\}.$$

It is easy to see that if p is constant, then $L^{p(\cdot)}(\mathbb{R}^n)$ is nothing but the standard Lebesgue space $L^p(\mathbb{R}^n)$. The space $L^{p(\cdot)}(\mathbb{R}^n)$ is referred to as a variable Lebesgue space.

We will always suppose that

$$1 < p_- := \operatorname*{ess\,inf}_{x\in\mathbb{R}^n} p(x), \quad \operatorname*{ess\,sup}_{x\in\mathbb{R}^n} p(x) =: p_+ < \infty. \tag{5.1}$$

Under these conditions, the space $L^{p(\cdot)}(\mathbb{R}^n)$ is separable and reflexive, and its associate space is isomorphic to $L^{p'(\cdot)}(\mathbb{R}^n)$, where

$$1/p(x) + 1/p'(x) = 1 \quad \text{for a.e.} \quad x \in \mathbb{R}^n$$

(see e.g. [**CF13**, Chap. 2] or [**DHHR11**, Chap. 3]).

5.2. The Hardy-Littlewood maximal function on variable Lebesgue spaces. By $\mathcal{B}_M(\mathbb{R}^n)$ denote the set of all measurable functions $p : \mathbb{R}^n \to [1,\infty]$ such that (5.1) holds and the Hardy-Littlewood maximal operator is bounded on the variable Lebesgue space $L^{p(\cdot)}(\mathbb{R}^n)$.

To provide a simple sufficient conditions guaranteeing that $p \in \mathcal{B}_M(\mathbb{R}^n)$, we need the following definition. Given a function $r : \mathbb{R}^n \to \mathbb{R}$, one says that r is locally log-Hölder continuous if there exists a constant $C_0 > 0$ such that

$$|r(x) - r(y)| \le \frac{C_0}{-\log|x-y|}$$

for all $x, y \in \mathbb{R}^n$ such that $|x-y| < 1/2$. One says that $r : \mathbb{R}^n \to \mathbb{R}$ is log-Hölder continuous at infinity if there exist constants C_∞ and r_∞ such that for all $x \in \mathbb{R}^n$,

$$|r(x) - r_\infty| \le \frac{C_\infty}{\log(e+|x|)}.$$

The class of functions $r : \mathbb{R}^n \to \mathbb{R}$ that are simultaneously locally log-Hölder continuous and log-Hölder continuous at infinity is denoted by $LH(\mathbb{R}^n)$. From [**CF13**, Proposition 2.3 and Theorem 3.16] we extract the following.

THEOREM 5.1. *Let $p \in LH(\mathbb{R}^n)$ satisfy (5.1). Then $p \in \mathcal{B}_M(\mathbb{R}^n)$.*

Although the latter result provides a nice sufficient condition for the boundedness of the Hardy-Littlewood maximal operator on the variable Lebesgue space $L^{p(\cdot)}(\mathbb{R}^n)$, it is not necessary. Notice that all functions in $LH(\mathbb{R}^n)$ are continuous and have limits at infinity. Lerner [**L05**] (see also [**CF13**, Example 4.68]) proved that if $p_0 > 1$ and $\mu \in \mathbb{R}$ is sufficiently close to zero, then the following variable exponent

$$p(x) = p_0 + \mu \sin(\log\log(1 + \max\{|x|, 1/|x|\})), \quad x \neq 0,$$

belongs to $\mathcal{B}_M(\mathbb{R})$. It is clear that the function p does not have limits at zero or infinity. We refer to the recent monographs [**CF13, DHHR11**] for further discussions concerning the fascinating and still mysterious class $\mathcal{B}_M(\mathbb{R}^n)$.

We will need the following remarkable result proved by Diening [**D05**, Theorem 8.1] (see also [**DHHR11**, Theorem 5.7.2] and [**CF13**, Corollary 4.64]).

THEOREM 5.2. *We have $p \in \mathcal{B}_M(\mathbb{R}^n)$ if and only if $p' \in \mathcal{B}_M(\mathbb{R}^n)$.*

5.3. Boundedness of maximally modulated Calderón-Zygmund singular integral operators on variable Lebesgue spaces. From Theorems 3.1 and 5.2 we immediately get the following.

COROLLARY 5.3. *Let $p \in \mathcal{B}_M(\mathbb{R}^n)$. Suppose T is a Calderón-Zygmund operator and $\Phi = \{\phi_\alpha\}_{\alpha\in\mathcal{A}}$ is a family of measurable real-valued functions indexed by an arbitrary set $\mathcal{A}$. If T^Φ is of weak type (r, r) for all $r \in (1, \infty)$, then T^Φ extends to a bounded operator on the variable Lebesgue space $L^{p(\cdot)}(\mathbb{R}^n)$.*

In turn, Corollary 5.3 and Lemma 3.2 yield the following.

COROLLARY 5.4. *If $p \in \mathcal{B}_M(\mathbb{R})$, then the maximally modulated Hilbert transform $\mathcal{C}$ extends to a bounded operator on the variable Lebesgue space $L^{p(\cdot)}(\mathbb{R})$.*

5.4. Boundedness of pseudodifferential operators with $L^\infty(\mathbb{R}, V(\mathbb{R}))$ symbols on variable Lebesgue spaces. Combining Lemma 4.2 with Theorem 5.2 we arrive at the following.

COROLLARY 5.5. *Suppose $p \in \mathcal{B}_M(\mathbb{R})$. Then the operator S_*, defined for the functions $f \in C_0^\infty(\mathbb{R})$ by (1.1), extends to a bounded operator on the variable Lebesgue space $L^{p(\cdot)}(\mathbb{R})$.*

From Theorems 4.3 and 5.2, taking into account Corollary 5.5, we get the following.

COROLLARY 5.6. *If $p \in \mathcal{B}_M(\mathbb{R})$ and $a \in L^\infty(\mathbb{R}, V(\mathbb{R}))$, then the pseudodifferential operator $a(x, D)$, defined for the functions $f \in C_0^\infty(\mathbb{R})$ by the iterated integral (1.3), extends to a bounded linear operator on the space $L^{p(\cdot)}(\mathbb{R})$ and*

$$\|a(x, D)\|_{\mathcal{B}(L^{p(\cdot)}(\mathbb{R}))} \leq 2\|S_*\|_{\mathcal{B}(L^{p(\cdot)}(\mathbb{R}))}\|a\|_{L^\infty(\mathbb{R},V(\mathbb{R}))}.$$

6. Compactness of pseudodifferential operators with non-regular symbols on variable Lebesgue spaces

6.1. Compactness of pseudodifferential operators with $L^\infty(\mathbb{R}, V(\mathbb{R}))$ symbols on standard Lebesgue spaces. We start with the case of constant exponents.

THEOREM 6.1 ([**K-Yu07**, Theorem 4.1]). *Let $1 < r < \infty$. If $a \in L^\infty(\mathbb{R}, V(\mathbb{R}))$ and*

(a) $a(x, \pm\infty) = 0$ *for almost all* $x \in \mathbb{R}$;
(b) $\lim\limits_{|x|\to\infty} V(a(x,\cdot)) = 0$;
(c) *for every* $N > 0$,

$$\lim_{L\to+\infty} \operatorname*{ess\,sup}_{|x|\le N} \left(V_{-\infty}^{-L}(a(x,\cdot)) + V_L^{+\infty}(a(x,\cdot))\right) = 0;$$

then the pseudodifferential operator $a(x, D)$ is compact on the standard Lebesgue space $L^r(\mathbb{R})$.

6.2. Transferring the compactness property from standard Lebesgue spaces to variable Lebesgue spaces. For a Banach space E, let $\mathcal{L}(E)$ and $\mathcal{K}(E)$ denote the Banach algebra of all bounded linear operators and its ideal of all compact operators on E, respectively.

THEOREM 6.2. *Let $p_j : \mathbb{R}^n \to [1,\infty]$, $j = 0, 1$, be a.e. finite measurable functions, and let $p_\theta : \mathbb{R}^n \to [1,\infty]$ be defined for $\theta \in [0,1]$ by*

$$\frac{1}{p_\theta(x)} = \frac{\theta}{p_0(x)} + \frac{1-\theta}{p_1(x)}, \quad x \in \mathbb{R}^n.$$

Suppose A is a linear operator defined on $L^{p_0(\cdot)}(\mathbb{R}^n) + L^{p_1(\cdot)}(\mathbb{R}^n)$.

(a) *If $A \in \mathcal{L}(L^{p_j(\cdot)}(\mathbb{R}^n))$ for $j = 0, 1$, then $A \in \mathcal{L}(L^{p_\theta(\cdot)}(\mathbb{R}^n))$ for all $\theta \in [0,1]$ and*

$$\|A\|_{\mathcal{L}(L^{p_\theta(\cdot)}(\mathbb{R}^n))} \le 4\|A\|^{\theta}_{\mathcal{L}(L^{p_0(\cdot)}(\mathbb{R}^n))}\|A\|^{1-\theta}_{\mathcal{L}(L^{p_1(\cdot)}(\mathbb{R}^n))}.$$

(b) *If $A \in \mathcal{K}(L^{p_0(\cdot)}(\mathbb{R}^n))$ and $A \in \mathcal{L}(L^{p_1(\cdot)}(\mathbb{R}^n))$, then $A \in \mathcal{K}(L^{p_\theta(\cdot)}(\mathbb{R}^n))$ for all $\theta \in (0,1)$.*

Part (a) is proved in [**DHHR11**, Corollary 7.1.4] under the more general assumption that p_j may take infinite values on sets of positive measure (and in the setting of arbitrary measure spaces). Part (b) follows from a general interpolation theorem by Cobos, Kühn, and Schonbeck [**CKS92**, Theorem 3.2] for the complex interpolation method for Banach lattices satisfying the Fatou property. Indeed, the complex interpolation space $[L^{p_0(\cdot)}(\mathbb{R}^n), L^{p_1(\cdot)}(\mathbb{R}^n)]_{1-\theta}$ is isomorphic to the variable Lebesgue space $L^{p_\theta(\cdot)}(\mathbb{R}^n)$ (see [**DHHR11**, Theorem 7.1.2]), and $L^{p_j(\cdot)}(\mathbb{R}^n)$ have the Fatou property (see [**DHHR11**, p. 77]).

The following characterization of the class $\mathcal{B}_M(\mathbb{R}^n)$ was communicated to the authors of [**KS13**] by Diening.

THEOREM 6.3 ([**KS13**, Theorem 4.1]). *If $p \in \mathcal{B}_M(\mathbb{R}^n)$, then there exist constants $p_0 \in (1,\infty)$, $\theta \in (0,1)$, and a variable exponent $p_1 \in \mathcal{B}_M(\mathbb{R}^n)$ such that*

$$\frac{1}{p(x)} = \frac{\theta}{p_0} + \frac{1-\theta}{p_1(x)}, \quad x \in \mathbb{R}^n. \tag{6.1}$$

From the above two theorems we obtain the following result, which allows us to transfer the compactness property from standard Lebesgue spaces to variable Lebesgue spaces.

LEMMA 6.4. *Let $A \in \mathcal{L}(L^{p(\cdot)}(\mathbb{R}^n))$ for all $p \in \mathcal{B}_M(\mathbb{R}^n)$. If $A \in \mathcal{K}(L^r(\mathbb{R}^n))$ for some $r \in (1,\infty)$, then $A \in \mathcal{K}(L^{p(\cdot)}(\mathbb{R}^n))$ for all $p \in \mathcal{B}_M(\mathbb{R}^n)$.*

PROOF. By the hypothesis, the operator A is bounded on all standard Lebesgue spaces $L^r(\mathbb{R}^n)$ with $1 < r < \infty$. From the classical Krasnosel'skii interpolation theorem (Theorem 6.2(b) with constant exponents) it follows that $A \in \mathcal{K}(L^r(\mathbb{R}^n))$ for all $1 < r < \infty$. If $p \in \mathcal{B}_M(\mathbb{R}^n)$, then in view of Theorem 5.2 there exist $p_0 \in (1, \infty)$, $\theta \in (0, 1)$, and a variable exponent $p_1 \in \mathcal{B}_M(\mathbb{R}^n)$ such that (6.1) holds. Since $A \in \mathcal{L}(L^{p_1(\cdot)}(\mathbb{R}^n))$ and $A \in \mathcal{K}(L^{p_0}(\mathbb{R}^n))$, from Theorem 6.2(b) we obtain $A \in \mathcal{K}(L^{p(\cdot)}(\mathbb{R}^n))$. □

6.3. Compactness of pseudodifferential operators with $L^\infty(\mathbb{R}, V(\mathbb{R}))$ symbols on variable Lebesgue spaces. Combining Corollary 5.6 and Theorem 6.1 with Lemma 6.4, we arrive at our last result.

COROLLARY 6.5. *Suppose $p \in \mathcal{B}_M(\mathbb{R})$. If $a \in L^\infty(\mathbb{R}, V(\mathbb{R}))$ satisfies the hypotheses* (a)–(c) *of Theorem 6.1, then the pseudodifferential operator $a(x, D)$ is compact on the variable Lebesgue space $L^{p(\cdot)}(\mathbb{R})$.*

References

[BS88] C. Bennett and R. Sharpley, *Interpolation of Operators*. Academic Press, New York, 1988.

[CKS92] F. Cobos, T. Kühn, and T. Schonbek, *One-sided compactness results for Aronszajn-Gagliardo functors.* J. Funct. Analysis **106** (1992), 274–313.

[CF13] D. Cruz-Uribe and A. Fiorenza, *Variable Lebesgue Spaces*. Birkhäuser, Basel, 2013.

[DPL13] F. Di Plinio and A. K. Lerner, *On weighted norm inequalities for the Carleson and Walsh-Carleson operators.* arXiv:1312.0833 (2013).

[D05] L. Diening, *Maximal function on Musielak-Orlicz spaces and generlaized Lebesgue spaces.* Bull. Sci. Math. **129** (2005), 657–700.

[DHHR11] L. Diening, P. Harjulehto, P. Hästö, and M. Růžička, *Lebesgue and Sobolev Spaces with Variable Exponents*. Lecture Notes in Mathematics **2017**, Springer, Berlin, 2011.

[D91] E. M. Dyn'kin, *Methods of the theory of singular integrals (the Hilbert transform and Calderon-Zygmund theory).* In "Commutative Harmonic Analysis I", Itogi Nauki i Tekhniki. Ser. Sovrem. Probl. Mat. Fund. Napr., **15**, VINITI, Moscow (1987), 197-292 (in Russian). English translation: Commutative harmonic analysis I. General survey. Classical aspects, Encycl. Math. Sci. **15** (1991), 167–259.

[FS72] Ch. Fefferman and E. M. Stein, *H^p spaces of several variables.* Acta Math. **129** (1972), 137–193.

[G09] L. Grafakos, *Modern Fourier Analysis.* 2nd ed. Graduate Texts in Mathematics **250**. New York, Springer, 2009.

[GMS05] L. Grafakos, J. M. Martell, and F. Soria, *Weighted norm inequalities for maximally modulated singular integral operators.* Math. Ann. **331** (2005), 359–394.

[K-A14] A. Yu. Karlovich, *Boundedness of pseudodifferential operators on Banach function spaces.* In: "Operator Theory, Operator Algebras and Applications". Operator Theory: Advances and Applications **242** (2014), 185–195.

[KS13] A. Yu. Karlovich and I. M. Spitkovsky, *Pseudodifferential operators on variable Lebesgue spaces.* In: "Operator Theory, Pseudo-Differential Equations, and Mathematical Physics. The Vladimir Rabinovich Anniversary Volume". Operator Theory: Advances and Applications **228** (2013), 173–183.

[KS14] A. Yu. Karlovich and I. M. Spitkovsky, *The Cauchy singular integral operator on weighted variable Lebesgue spaces.* In: "Concrete Operators, Spectral Theory, Operators in Harmonic Analysis and Approximation". Operator Theory: Advances and Applications **236** (2014), 275–291.

[K-Yu07] Yu. I. Karlovich, *Algebras of pseudo-differential operators with discontinuous symbols.* In: "Modern Trends in Pseudo-Differential Operators". Operator Theory: Advances and Applications **172** (2007), 207–233.

[K-Yu12] Yu. I. Karlovich, *Boundedness and compactness of pseudodifferential operators with non-regular symbols on weighted Lebesgue spaces.* Integr. Equ. Oper. Theor. **73** (2012), 217–254.

[L05] A. K. Lerner, *Some remarks on the Hardy-Littlewood maximal function on variable L^p spaces*. Math. Z. **251** (2005), 509-521.

[L10] A. K. Lerner, *Some remarks on the Fefferman-Stein inequality*. J. Anal. Math. **112** (2010), 329-349.

[LO10] A. K. Lerner and S. Ombrosi, *A boundedness criterion for general maximal operators*. Publ. Mat. **54** (2010), 53-71.

[LP07] A. K. Lerner and C. Pérez, *A new characterization of the Muckenhoupt A_p weights through an extension of the Lorentz-Shimogaki theorem*. Indiana Univ. Math. J. **56** (2007), 2697-2722.

[MS13] C. Muscalu and W. Schlag, *Classical and Multilinear Harmonic Analysis*. Vol. II. Cambridge Studies in Advanced Mathematics **138**. Cambridge University Press, Cambridge, 2013.

[N55] I. P. Natanson, *Theory of Functions of a Real Variable*. Frederick Ungar Publishing Co., New York, 1955.

[S93] E. Stein, *Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals*. Princeton Uinversity Press, Princeton, NJ, 1993.

[S67] L. Schwartz, *Analyse Mathématique. Cours I*. Hermann, Paris, 1967.

Centro de Matemática e Aplicações (CMA) and Departamento de Matemática, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, Quinta da Torre, 2829–516 Caparica, Portugal

E-mail address: oyk@fct.unl.pt